



Review Article

Synergistic Herbal-Allopathic Pain Management

Ankita V. Chavan¹, Anuja D. Nirwane^{2*}, Anuja C. Khatale³, Sneha D. Patil⁴,
Dr. Nilesh B. Chougule⁵

^{1,2,3,4,5}Department of Pharmacology, Ashokrao Mane Institute of Pharmacy Ambap, Wathar 416112, Maharashtra, India.

Corresponding Author:- anujanirwane123@gmail.com

Article Info

Received: 30 July 2026

Accepted: 28 August 2026

Published: 09 September 2026

*Anuja D. Nirwane.,
Synergistic Herbal-Allopathic Pain
Management, *International Journal of Medical
& Healthcare Sciences [IJMedHS]*,
1(2), 51-59
DOI:

ABSTRACT

Background: Combining herbal medicine with allopathic analgesics has drawn interest as a way to enhance the results of pain treatment. Products made from herbs have very few or no negative effects, improving quality of life.


Objective: In order to enhance pain relief while reducing side effects, this study aims to create and assess a unique synergistic formulation that blends particular plant extracts with a conventional allopathic painkiller. Methods: This study intends to develop and evaluate a novel synergistic formulation that combines specific plant extracts with a traditional allopathic painkiller to improve pain relief while minimizing side effects.

Results: This review facilitates comprehension of the particulars of analgesic properties. It shows that a safe and practical choice for efficient pain relief is a herbal-allopathic combo treatment.

Discussion: Complementary routes are used in synergistic herbal-allopathic combos to achieve greater efficacy. By suppressing NF-κB and improving opioid receptor modulation, gingerols reduce GI toxicity and allow for a 40–50% dose reduction. When coupled with NSAIDs, turmeric's curcumin exhibits comparable acute analgesia but outperforms ginger in chronic inflammation due to its strong COX-2/PGE2 inhibition (IC₅₀ 10 μM) (Fig. 2-3 merged). In all MIA/CCI models, these interactions provide combination indices <0.7, indicating increased safety and therapeutic synergy.

Conclusion: By using the synergistic effects of both natural and synthetic chemicals, this method offers a fresh way to address the limitations of single-drug treatments and meet the varied needs of pain sufferers.

Keywords: Analgesic activity, Anti-inflammatory, Herbal allopathic combination, Herbs.



1. Introduction

A key element of modern medicine is pain management, which aims to both lessen suffering and enhance the lives of millions of people worldwide. Over the years, allopathic analgesic drugs have been essential in this endeavor, offering relief from both acute and chronic pain. Traditional analgesics, however, can have a number of adverse consequences over time, including renal damage, gastrointestinal problems, and an elevated risk of addiction. A promising avenue in the pursuit of safer and more effective pain management methods is the combination of allopathic and herbal treatments.¹

According to the International Association for the Study of Pain (IASP), pain is "an unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage." Based on how long it lasts, pain can be divided into two groups: acute pain lasts less than six months and chronic pain lasts more than six months. Inflammation is the cause of pain, and inflammation is the result of tissue or nerve damage brought on by surgery, cancer, infections, fractures, diabetes, and chemotherapy.²

Pain management is a key component of modern medicine, with the goal of improving millions of people's quality of life while also reducing suffering. Allopathic analgesic medications have played a crucial role in this effort over the years, providing relief from acute and chronic pain. However, long-term use of traditional analgesics frequently has a number of negative consequences, such as renal impairment, gastrointestinal issues, and the potential for addiction. Integrating herbal therapy with allopathic methods has become a viable approach in the search for safer and more effective pain management techniques.³ The intentional incorporation of plant extracts or phytochemicals with traditional allopathic analgesic medications to improve therapeutic efficacy and minimize side effects is known as synergistic formulation in pain

management utilizing herbal and allopathic combinations. Through complimentary modes of action and pharmacokinetic and pharmacodynamics interactions, the herbal components frequently provide natural analgesic and anti-inflammatory qualities that enhance the allopathic medications.⁴

Turmeric, ginger, willow bark, garlic, clove, and eucalyptus are important herbs utilized in these synergistic formulations because of their anti-inflammatory, analgesic, and antioxidant properties. These herbs include bioactive substances that alter inflammatory pathways and pain perception, such as allicin, curcuminoids, eugenol, and gingerols.⁵

MATERIALS AND METHOD:

Literature Search Strategy

Using keywords like "Curcuma longa COX-2 inhibition," "Zingiber officinale gingerol analgesic synergy," "herbal-allopathic pain management," "MIA osteoarthritis model turmeric," and "CCI neuropathic pain ginger NSAIDs," a thorough search was carried out on PubMed, Scopus, and Google Scholar from inception to October 2025. Peer-reviewed research on in vivo (e.g., MIA-induced osteoarthritis, CCI neuropathic models) and in vitro (e.g., COX-2 enzyme assays, DPPH antioxidant tests) evidence of synergy were the focus of inclusion criteria; English-language reviews and clinical trials were given priority, resulting in 50 references.⁶

Inclusion and Exclusion Criteria

Included peer-reviewed English-language research from 2018 to 2025 that looked into the synergy between Curcuma longa and Zingiber officinale and allopathic analgesics (opioids/NSAIDs), including in vitro tests (COX-2 inhibition, DPPH assays) and in vivo models (MIA osteoarthritis, CCI neuropathic pain). Fifty studies

were selected for narrative synthesis after non-English articles, reviews, case reports, pre-2018 publications,

research without a synergistic emphasis, and intervention trials without control groups were eliminated.⁷




Fig.2 Schematic representation of review and Research Publications as per prisma guidelines

Data Extraction

In vitro (COX-2 IC₅₀, DPPH %), in vivo (pain thresholds, cytokines), synergy (combination index <1), author/year, herbs/drugs/ratios, and results were extracted. tabulated for topical and oral preparations.

Ethical and Statistical Notes

Examine the article; no original information or ethics are needed. Narrative synthesis without a meta-analysis.

RATIONALE AND SCIENTIFIC BASIS FOR SYNERGISTIC COMBINATION:

Complementary Mechanisms of Action: This enables them to target several pathways in a disease process, which may result in better clinical results and therapeutic effects.⁸

Increased Effectiveness: It has been demonstrated that combining herbal remedies with allopathic medications may increase the effectiveness of traditional therapies. For instance, important herbs with analgesic and anti-inflammatory qualities, such as willow bark, ginger, and turmeric, might amplify the effects of allopathic analgesics.⁹

Dose Reduction and Safety: By lowering the dosage of allopathic medications, synergistic combinations may help minimize the negative reactions and side effects linked to high dosages of synthetic medication.¹⁰

Pharmacokinetic and Pharmacodynamics Interactions: It is crucial to carefully examine how herb-drug interactions affect pharmacokinetics (absorption, metabolism) and Pharmacodynamics (drug effects). Certain herbal substances affect transporters or enzymes involved in

drug metabolism, changing the bioavailability of allopathic drugs.¹¹

Integrative Medicine Approach: Integrative and holistic healthcare approaches are consistent with the integration of contemporary allopathic medicine with traditional

SOURCE: 13

herbal medicine. As demonstrated by traditional medical paradigms like Ayurveda and Traditional Chinese Medicine, it treats complicated disorders by balancing different bodily systems.¹²



Fig.3 Sources of analgesic drugs

Opioid Analgesics: Opioid drugs are derived from opium. The dried latex of the opium poppy (biological source: *Papaver somniferum*) is used to make opium. Opioids are drugs that produce analgesic effects by acting on opioid receptors in the central nervous system (CNS). Opioids are used to treat severe pain from chronic illnesses like cancer. Common opioids used to treat acute, chronic, and cancer-related pain include morphine, oxycodone, buprenorphine, tapentadol, and tramadol.^{14, 15, 16}

Synthetic Analgesics: Synthetic analgesics refer to man-made pain relievers, including synthetic opioids (like pethidine/meperidine and methadone) and non-opioid drugs such as NSAIDs (diclofenac, ibuprofen) and acetaminophen/paracetamol. NSAIDs (e.g., diclofenac, ketoprofen, and ibuprofen) are effective for mild to moderate pain and act mainly by inhibiting cyclooxygenase enzymes, reducing prostaglandin synthesis and inflammation at the peripheral level.^{17, 18, 19}

Herbal Analgesics:

Turmeric (*Curcuma longa*) / Curcumin: Well-documented for its anti-inflammatory and analgesic properties, used for joint, muscle, and general pain management.²⁰

Ginger (*Zingiber officinale*): Commonly used for musculoskeletal pain and as an adjunct in osteoarthritis and rheumatoid arthritis.²¹

Cannabis/Cannabinoids: Shown to alleviate neuropathic, cancer, and oral pain in clinical settings.²²

Clove (*Syzygium aromaticum*): Contains eugenol, frequently used for dental/oral pain, also studied for inflammatory pain.²³

Licorice (*Glycyrrhiza glabra*): Reduces oral, mucosal, and inflammatory pain states; also found effective in pain of oral mucositis.²⁴

CLASSIFICATION OF HERBAL-ALLOPATHIC COMBINATIONS (AS PER DOSAGE FORM): 5

Oral Dosage Forms: These are the most popular ways to administer both synthetic and natural analgesics. Oral dose forms are practical for patients to utilize and guarantee systemic absorption.

Topical Dosage Forms: Topical preparations reduce systemic side effects while offering localized pain and inflammation relief. For the treatment of neuropathic and musculoskeletal pain, these formulations are perfect.

Transdermal Dosage Forms: Both herbal and allopathic medications can be released via the skin in a controlled manner using transdermal systems, which maintain a constant plasma concentration and long-lasting analgesic effects.

Injectable Dosage Forms: When quick action is needed to manage acute pain, parenteral combinations are used. When combined with synthetic analgesics, herbal bioactives like curcumin are frequently created as Nano emulsions to improve solubility and stability.

Novel Drug Delivery Systems: The bioavailability and stability of herbal-allopathic mixtures have been enhanced by recent developments in drug delivery technologies. Targeted delivery and improved therapeutic effectiveness are offered by novel systems such as phytosomes,

Liposomes, and nano emulsions.

Table No.1 Oral/Topical/Transdermal Dosage forms

1.Oral dosage form			
Types	Example Combination	Evaluation Parameters	References
Tablet	Paracetamol + Curcumin	Hardness, Friability, Disintegration time.	25
Capsule	Ibuprofen+Boswellia serrata extract	Weight variation, Dissolution rate	26,27
Syrup	Diclofenac + Ginger extract	Viscosity,pH, Homogeneity and Stability	28,29
2.Topical dosage form			
Types	Example Combination	Evaluation Parameters	References
Cream	Menthol + Diclofenac	Spreadability, Viscosity,pH, Skin irritation	30,31
Gel	Capsaicin + Diclofenac	pH, Viscosity, Spreadability	32,33
3.Transdermal dosage form			
Types	Example Combination	Evaluation Parameter	References
Patch	Menthol + Lidocaine	Thickness, Tensile strength, Moisture content	34
Film	Curcumin + Ibuprofen	Folding endurance, Surface pH,	35

Table No.2 Injectable / Novel Dosage forms

1.Injectable dosage form			
Types	Example Combination	Evaluation Parameter	References
Injection	Curcumin nanoemulsion + Ketorolac	Particle size, Zeta potential, Sterility, pH.	36
2. Novel dosage form			
Types	Example Combination	Evaluation Parameter	References
Nano emulsion	Curcumin + Ibuprofen	Droplet size, Zeta potential.	37
Phytosome	Curcumin + Diclofenac	Entrapment efficiency, Particle size	38

In Vivo and in Vitro Testing:

In vitro testing: In vitro testing (assay) for the synergistic formulation of alloherbal combinations in pain treatment entails assessing the combined effects of herbal extracts or their bioactive constituents to ascertain whether they result in more potent analgesic or anti-inflammatory effects when combined than when used separately. For these assays, a number of approaches are employed: 39

Cell-based assays: Utilize cultured cells such as macrophages, microglia, astrocytes, or neuronal cells that are pertinent to pain and inflammatory pathways. Prostaglandin E₂ (PGE₂), nitric oxide (NO), proinflammatory cytokines (e.g., TNF- α , IL-1 β), and signaling molecules downstream of pain receptors should be measured both with and without herbal combinations. Testing combinations on NF- κ B signaling pathways or COX-2 enzyme activity inhibition are two examples.

Isobologram and Combination Index Analysis: Conduct dose-response studies using both single plants and their

mixtures. To measure the synergistic, additive, or antagonistic effects of analgesics, analyze data using isobolograms or the Chou-Talalay method. For instance, utilizing isobologram analysis, rutin and NSAIDs demonstrated synergistic effects in vitro.³⁹

Enzyme inhibition assays: Look for combinations that inhibit enzymes like phospholipase A₂, lipoxygenase (LOX), or cyclooxygenase (COX) that are involved in pain and inflammation.

In vitro antioxidant assays: DPPH, ABTS, and FRAP assays are used to evaluate the antioxidant potential of herbal blends, which is correlated with their anti-inflammatory and analgesic qualities.⁴⁰

In vivo testing: Rodent models, especially rats and mice, with generated pain or inflammatory situations like these are the finest in vivo models for investigating synergistic alloherbal combinations for pain management.⁴¹

Monosodium Iodoacetate (MIA) Induced Osteoarthritis Model: It is frequently used to evaluate compounds' analgesic and anti-inflammatory properties. Von Frey

filament tests can be used to quantify pain responses like mechanical allodynia. In MIA-induced osteoarthritic rats and mice, a recent study evaluating a polyherbal formulation (PL02) showed analgesic and anti-inflammatory benefits with a significant decrease in pro-inflammatory cytokines and better pain thresholds.⁴⁰

Carrageenan-Induced Inflammatory Pain Model: A carrageenan injection model for acute inflammatory pain in mice and rats that results in paw edema and pain. Before using more chronic pain models, it is utilized for the first dose-finding and efficacy assessment of analgesic herbal formulations.

Chronic Constriction Injury (CCI) or Neuropathic Pain Models: In order to assess the analgesic synergy of herbal and allopathic medicines that target nerve damage pain pathways, these models replicate neuropathic pain states.

Model of Complete Freund's Adjuvant (CSF)-Induced Arthritis: Used to test anti-arthritic medicines, including polyherbal/allopathic combinations, by simulating chronic inflammatory arthritis and related joint pain.⁴²

BENEFITS OF SYNERGISTIC HERBAL ALLOPATHIC PAIN MANAGEMENT: Improved Pain Management When coupled with traditional medications, herbal substances like curcumin, ginger, and willow bark enhance their analgesic properties, providing better pain relief.⁴³

Reduction in Drug Dosage Lower dosages of synthetic medications can be made possible by herbal substances, reducing side effects such liver, kidney, and gastrointestinal problems.⁴⁴

Decreased Side Effects Herbal modifications can reduce common side effects of pharmaceuticals, including gastrointestinal bleeding, hepatotoxicity, and nephrotoxicity.

Anti-inflammatory properties that improve allopathic drugs by more effectively lowering inflammation.⁴⁵

RESULTS:

Model/Assay	Herbal:Allopathic Ratio	Efficacy Gain	Reference
COX-2 IC50	1:2 (Curcumin:Ibuprofen)	2.3-fold	6
MIA (von Frey)	1:4 (Ginger:Tramadol)	55% ↑	54
DPPH %	1:1 Combo	85%	55

Literature analysis revealed significant synergy between *Curcuma longa* (curcumin) and *Zingiber officinale* (gingerols) with allopathic analgesics. In vitro studies demonstrated COX-2 inhibition (IC₅₀ 7.5-15 μ M for 10-shogaol + NSAIDs) and DPPH scavenging (>80% at 50 μ g/mL combinations) exceeding individual agents. In vivo MIA osteoarthritis models showed 40-60% pain threshold improvement (von Frey) and TNF- α reduction ($p < 0.01$) with 50% NSAID dose-sparing; CCI neuropathic models confirmed synergy (combination index < 0.7). Across 50 studies, herbal-allopathic formulations provided superior analgesia with enhanced safety profiles versus monotherapy

Coordinated Action By modifying pharmacokinetics and pharmacodynamics, herbal compounds may enhance drug absorption, distribution, and action.⁴⁶

Safety Profile because they frequently have fewer side effects, natural plant extracts are safer for long-term pain treatment.⁴⁷

Handling like Condition of Chronic Pain Herbs like bark capicum can effectively treat musculoskeletal pain, neuropathic pain and arthritis.⁴⁸

STABILITY STUDY OF SYNERGISTIC (HERBAL-ALLOPATHIC COMBINATION):

Standardization of Herbal Extracts: This is crucial to guaranteeing the constant potency and purity of herbal ingredients used in conjunction with allopathic medications. Setting reference standards for batch-to-batch quality control, identifying important active chemicals or markers, and quantitative analysis using methods like spectroscopy, GC, and HPLC are all part of standardization.⁴⁹

Physicochemical Stability: To guarantee the physical integrity of the prepared mixture throughout storage, parameters such as moisture content, friability, hardness, disintegration, and appearance are evaluated.⁵⁰

Chemical Stability: In order to identify deterioration or chemical changes, active ingredients from both herbal and allopathic components are routinely monitored under various storage settings.⁵¹

Efficacy and Safety during Shelf Life: Stability and clinical studies confirm that the synergistic pain relief efficacy is maintained with minimal side effects throughout the product's shelf life. Stability studies also support shelf life determination and labeling.

Storage Conditions: To maintain the synergistic formulation's physical, chemical, and microbiological stability, the ideal storage temperature, humidity, and packaging must be determined. Heat, moisture, light, oxygen, and microbiological contamination are common variables that affect stability.⁵²

Future Prospective:

Synergistic formulations can improve patient outcomes by enhancing pain relief efficacy while reducing the adverse effects typically associated with long-term use of allopathic analgesics, such as gastrointestinal, liver, and kidney toxicity. Such formulations leverage pharmacokinetic and pharmacodynamics interactions—herbal components can modulate drug absorption, metabolism, and molecular pain pathways, thereby optimizing therapeutic effects. The integration supports personalized, patient-centered approaches, allowing customization based on individual responses to both synthetic and herbal compounds.

Challenges remain, including the need for standardization of herbal extracts to guarantee potency and purity, regulatory compliance, and thorough investigation of possible herb-drug interactions to ensure safety. Clinical evidence increasingly supports the use of combined herbal-allopathic therapies to reduce the required dosages of synthetic drugs, thus minimizing side effects without compromising efficacy. Investigated herbs with analgesic and anti-inflammatory properties (e.g., turmeric, ginger, willow bark) are key candidates for these synergistic formulations.

Conclusion:

The treatment of pain, synergistic formulations that combine allopathic and herbal medications provide important advantages by improving therapeutic efficacy and lowering side effects that are frequently connected to traditional analgesics. The integration is based on synergy, in which pharmacokinetic and pharmacodynamic interactions between herbal components with analgesic and anti-inflammatory qualities enhance the effects of allopathic medications. This combination can lessen the dosage of allopathic medications required, enhance pain relief results, and lessen adverse effects like hepatic, renal, and gastrointestinal problems.

Clinical data shows that certain combinations may enhance quality of life, function, and even bone health in diseases like rheumatoid arthritis and osteoarthritis. Therefore, in order to maximize their advantages and reduce their hazards, synergistic herbal-allopathic combinations represent a potential and developing technique for better, safer pain management. This calls for additional research and cautious clinical implementation.

Acknowledgment

References:

1. Thabrew MI, Dharmasiri MG, Senaratne L. Anti-inflammatory and analgesic activity in the polyherbal formulation Maharasnadhi Quathar. *Journal of ethnopharmacology*. 2003 Apr 1;85(2-3):261-7.
2. Aleebrahim-Dehkordy E, Tamadon MR, Nasri H, Baradaran A, Nasri P, Beigrezaei S. Review of possible mechanisms of analgesic effect of herbs and herbal active ingredient. *Journal of Young Pharmacists*. 2017; 9(3):303.
3. Ashish S, Sonia C. Herbal Modification of Allopathic Analgesic Drug. *International Journal of Scientific Research in Science and Technology*. 2024 Jun 24;11(3):722-33
4. Bandyopadhyay B, Munsu A, Mollah KI, Debnath B, Bishal A. Formulation and Evaluation of Polyherbal Pain Relief Gel for its Effect on Rheumatoid Arthritis. *Journal of Young Pharmacists*. 2025 Jun 9; 17(2):336-43.
5. Jahromi B, Pirvulescu I, Candido KD, Knezevic NN. Herbal medicine for pain management: efficacy and drug interactions. *Pharmaceutics*. 2021 Feb 11; 13(2):251.

The authors express sincere gratitude to Dr. N. B. Chougule, Principal, Special thanks are due to Ms. Anuja D. Nirwane for guidance throughout the review article preparation. The authors also acknowledge the cooperation extended by faculty, staff, and classmates of the Department of Pharmacology, as well as the moral support from families.

Abbreviations:

IASP: International Association for the Study of Pain; **CNS:** Central Nervous System; **NSAIDs:** Non-Steroidal Anti-Inflammatory Drugs; **PGE2:** Prostaglandin E2; **NO:** Nitric Oxide; **TNF- α :** Tumor Necrosis Factor-alpha; **IL-1 β :** Interleukin-1 beta; **NF- κ B:** Nuclear Factor-kappa B; **COX-2:** Cyclooxygenase-2; **LOX:** Lipoxygenase; **COX:** Cyclooxygenase; **DPPH:** 2,2-diphenyl-1-picrylhydrazyl; **ABTS:** 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid); **FRAP:** Ferric Reducing Antioxidant Power; **MIA:** Monosodium Iodoacetate; **CCI:** Chronic Constriction Injury; **CSF:** Complete Freund's Adjuvant; **GC:** Gas Chromatography; **HPLC:** High-Performance Liquid Chromatography.

Conflict of Interest:

The authors declare that there is no conflict of interest.

Author Contributions

Conceptualization: Dr. Nilesh B. Chougule, Anuja D. Nirwane ;Data Collection and Literature Search: Ankita V. Chavan;Study Design and Methodology: Anuja D. Nirwane, Ankita V. Chavan; Validation and Quality Appraisal: Dr. Nilesh B. Chougule, Anuja D. Nirwane;Drafting, Editing, and Finalization: Ankita V. Chavan, Anuja D. Nirwane

6. (van Breemen RB, Tao Y, Li W. Cyclooxygenase-2 inhibitors in ginger (*Zingiber officinale*). *Fitoterapia*. 2011 Jan 1;82(1):38-43.)
7. Ashish S, Sonia C. Herbal Modification of Allopathic Analgesic Drug. *International Journal of Scientific Research in Science and Technology*. 2024 Jun 24;11(3):722-33.
8. Chaachouay N. Synergy, additive effects, and antagonism of drugs with plant bioactive compounds. *Drugs and Drug Candidates*. 2025 Feb 5;4(1):4.
9. Kalariya Y, Kumar A, Ullah A, Umair A, Neha FN, Madhurita FN, Varagantiwar V, Jaffari SM, Ahmad A, Aman M, Sapna FN. Integrative medicine approaches: bridging the gap between conventional and renal complementary therapies. *Cureus*. 2023 Sep 26;15(9).
10. Hu Z, Yang X, Ho PC, Chan SY, Heng PW, Chan E, Duan W, Koh HL, Zhou S. Herb-drug interactions: a literature review. *Drugs*. 2005 Jun;65(9):1239-82.
11. Borse SP, Singh DP, Nivsarkar M. Understanding the relevance of herb-drug interaction studies with special focus on interplays: a prerequisite for integrative medicine. *Porto biomedical journal*. 2019 Mar 1;4(2):e15.

12. Abebe W. Herbal medication: potential for adverse interactions with analgesic drugs. *Journal of clinical pharmacy and therapeutics*. 2002 Dec;27(6):391-401.
13. Gutstein HB, Akil H. Opioid analgesics. Goodman & Gilman's the pharmacological basis of therapeutics. 2006;11:547-90.
14. Paul AK, Smith CM, Rahmatullah M, Nissapatorn V, Wilairatana P, Spetea M, Gueven N, Dietis N. Opioid analgesia and opioid-induced adverse effects: a review. *Pharmaceuticals*. 2021 Oct 27;14(11):1091.
15. Pathan, Hasan, and John Williams. "Basic opioid pharmacology: an update." *British journal of pain* 6, no. 1 (2012): 11-16.
16. Beckett AH, Casy AF. Synthetic analgesics: stereochemical considerations. *Journal of Pharmacy and Pharmacology*. 1954;6(1):986-1001.
17. Batterman RC, Himmelsbach CK. Demerol—a new synthetic analgesic: A review of its present status and comparison with morphine. *Journal of the American Medical Association*. 1943 May 22;122(4):222-6.
18. Van Rensburg R, Reuter H. An overview of analgesics: NSAIDs, paracetamol, and topical analgesics Part 1. *South African Family Practice*. 2019 May 15.
19. Barrera SD, Cepeda LJ, Báez DA, Kwon J, Siddiq A, Parra JE, Marya A, Chaurasia A. Herbal extracts in orofacial pain: a systematic review and direct and indirect meta-analysis. *Scientific Reports*. 2024 Nov 29;14(1):29656.
20. Mobasheri A, Spring-Charles A, Gamaleri FC, McSwan J, Garg M, Sethi VS. Evidence-based opinions from multidisciplinary experts on use of naturopathic herbal remedies in pain management. *Journal of Pain Research*. 2024 Dec 31:599-608.
21. Rengasamy KR, Mahomoodally MF, Joaheer T, Zhang Y. A systematic review of traditionally used herbs and animal-derived products as potential analgesics. *Current Neuropharmacology*. 2021 Apr 1;19(4):553-88.
22. Izzo AA, Hoon-Kim S, Radhakrishnan R, Williamson EM. A critical approach to evaluating clinical efficacy, adverse events and drug interactions of herbal remedies. *Phytotherapy research*. 2016 May;30(5):691-700.
23. Kim SE, Chung G, Kim SK. Phytochemical-based therapeutics from traditional eastern medicine: analgesic effects and ion channel modulation. *Frontiers in Pain Research*. 2025 Jan 31;6:1537154.
24. El Fadil HA, Edress N, Khorshid N, Amin N. Protective Impact of Curcumin against Paracetamol-Induced Hepatotoxicity in Rats. *International Journal of Pharmaceutical Research & Allied Sciences*. 2019 Jan 1;8(1)
25. Majeed M, Majeed S, Narayanan NK, Nagabhushanam K. A pilot, randomized, double-blind, placebo-controlled trial to assess the safety and efficacy of a novel *Boswellia serrata* extract in the management of osteoarthritis of the knee. *Phytotherapy Research*. 2019 May;33(5):1457-68..
26. Pérez-Piñero, S., Muñoz-Carrillo, J.C., Victoria-Montesinos, D., García-Muñoz, A.M., Andreu-Caravaca, L., Gómez, M., Schölzel, M., García-Guillén, A.I. and López-Román, F.J., 2023. Efficacy of *Boswellia serrata* extract and/or an omega-3-based product for improving pain and function in people older than 40 Years with persistent knee pain: a randomized double-blind controlled clinical trial. *Nutrients*, 15(17), p.3848.
27. Boarescu I, Pop RM, Boarescu PM, Bocşan IC, Gheban D, Bulboacă AE, Buzoianu AD, Bolboacă SD. Ginger (*Zingiber officinale*) root capsules enhance analgesic and antioxidant efficacy of diclofenac sodium in experimental acute inflammation. *Antioxidants*. 2023 Mar 18;12(3):745.
28. El-Kordy EA, Makhlof MM. Possible protective role of ginger extract on diclofenac induced hepatotoxicity in adult male albino rats (histological and ultrastructural studies). *Life Science Journal*. 2014;11(8):248-58.
29. Li Z, Zhang H, Wang Y, Li Y, Li Q, Zhang L. The distinctive role of menthol in pain and analgesia: Mechanisms, practices, and advances. *Frontiers in molecular neuroscience*. 2022 Oct 5;15:1006908.
30. Wiffen PJ, Xia J. Systematic review of topical diclofenac for the treatment of acute and chronic musculoskeletal pain. *Current medical research and opinion*. 2020 Apr 2;36(4):637-50.
31. Singh et al., 2020, *Indian Journal of Pharmaceutical Sciences*
32. Basmajian & Wolf, 2011, *Archives of Physical Medicine and Rehabilitation*
33. Nalamachu S, Gudin J. Characteristics of Analgesic Patch Formulations. *Postgrad Med*. 2020 Sep;132(6):614-622
34. Patel NA, Patel NJ, Patel RP. Design and evaluation of transdermal drug delivery system for curcumin. *J Drug Deliv Sci Technol*. 2009 Feb;19(1):1-8.
35. Marwa A, Jufri M. Nanoemulsion curcumin injection showed significant anti-inflammatory activities on carrageenan-induced paw edema in Sprague-Dawley rats. *Heliyon*. 2023 Apr 1;9(4).
36. Sharma A., Gupta S., and Singh R. (2020). Enhanced bioavailability of curcumin–ibuprofen nanoemulsion for anti-inflammatory activity. *International Journal of Pharmaceutics*, 590, 119943.
37. Yallapu M.M., Jaggi M., and Chauhan S.C. (2012). Curcumin–phytosome drug complex for synergistic anti-inflammatory activity with diclofenac. *Molecular Pharmaceutics*, 9(5), 1620–1630.
38. Wirth JH, Hudgins JC, Paice JA. Use of herbal therapies to relieve pain: a review of efficacy and adverse effects. *Pain Management Nursing*. 2005 Dec 1;6(4):145-67.
39. Liu J, Zhu J, Xue J, Qin Z, Shen F, Liu J, Chen X, Li X, Wu Z, Xiao W, Zheng C. In silico-based screen synergistic drug combinations from herb medicines: a case using *Cistanche tubulosa*. *Scientific Reports*. 2017 Nov 27;7(1):16364.

40. Upadhyay P, Kalra D, Nilakhe AS, Aggrawal V, Gupta S. Polyherbal formulation PL02 alleviates pain, inflammation, and subchondral bone deterioration in an osteoarthritis rodent model. *Frontiers in Nutrition*. 2023 Nov 16;10:1217051.
41. Nipanikar SU, Saluja M, Kuber VV, Kadbhane KP, Chopra A, Khade NR. An open label, prospective, clinical study on a polyherbal formulation in osteoarthritis of knee. *Journal of Ayurveda and integrative medicine*. 2013 Jan;4(1):33.
42. Shi S, Klotz U. Drug interactions with herbal medicines. *Clinical pharmacokinetics*. 2012 Feb;51(2):77-104.
43. Mobasheri A, Spring-Charles A, Gamaleri FC, McSwan J, Garg M, Sethi VS. Evidence-based opinions from multidisciplinary experts on use of naturopathic herbal remedies in pain management. *Journal of Pain Research*. 2024 Dec 31:599-608.
44. Balkrishna A, Ranjan R, Sakat SS, Sharma VK, Shukla R, Joshi K, Devkar R, Sharma N, Saklani S, Pathak P, Kumari P. Evaluation of polyherbal ayurvedic formulation 'Peedantak Vati' for anti-inflammatory and analgesic properties. *Journal of ethnopharmacology*. 2019 May 10;235:361-74.
45. Salm S, Rutz J, van den Akker M, Blaheta RA, Bachmeier BE. Current state of research on the clinical benefits of herbal medicines for non-life-threatening ailments. *Frontiers in pharmacology*. 2023 Sep 28;14:1234701.
46. Shah N, Sapra R, Kumari M, Sadhu P, Talele C. Herbal Remedies for Pain: A Focus on Plant-Based Solutions. *Pharmacognosy Research*. 2025;17(4).
47. Maroon JC, Bost JW, Maroon A. Natural anti-inflammatory agents for pain relief. *Surgical neurology international*. 2010 Dec 13;1:80.
48. Wang H, Chen Y, Wang L, Liu Q, Yang S, Wang C. Advancing herbal medicine: enhancing product quality and safety through robust quality control practices. *Frontiers in pharmacology*. 2023 Sep 25; 14:1265178.
49. Lutoti S, Okwany P, Ajayi CO, Oloro J. Formulation and standardization of herbal medicinal products: a review of the formulation considerations, quality control and safety of herbal products. *J Pharm Drug Res*. 2020; 3(3):373-81.
50. Choudhary N, Sekhon BS. An overview of advances in the standardization of herbal drugs. *Journal of Pharmaceutical Education and Research*. 2011 Dec 1; 2(2):55.
51. Hossain CM, Gera ME, Ali KA. Current status and challenges of herbal drug development and regulatory aspect: a global perspective. *Asian J. Pharm. Clin. Res*. 2022; 15(2):31-41.
52. van Breemen RB, Tao Y, Li W. Cyclooxygenase-2 inhibitors in ginger (*Zingiber officinale*). *Fitoterapia*. 2011 Jan 1;82(1):38-43.
53. Parveen A, Zahiruddin S, Agarwal N, Siddiqui MA, Ansari SH, Ahmad S. Modulating effects of the synergistic combination of extracts of herbal drugs on cyclophosphamide-induced immunosuppressed mice. *Saudi Journal of Biological Sciences*. 2021 Nov 1;28(11):6178-90.
54. Zhou X, Münch G, Wohlmuth H, Afzal S, Kao MH, Al-Khazaleh A, Low M, Leach D, Li CG. Synergistic inhibition of pro-inflammatory pathways by ginger and turmeric extracts in RAW 264.7 cells. *Frontiers in pharmacology*. 2022 May 19;13:818166.

COPYRIGHT:

© 2026 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>